

SUSY: From the Basics to Collider Phenomenology

Sven Heinemeyer, IFCA (Santander)

Louvain, 05/2007

1. Introduction to SUSY
2. SUSY Lagrangians and the MSSM
3. “Simplified versions” and special sectors
4. SUSY Phenomenology

SUSY: From the Basics to Collider Phenomenology

Sven Heinemeyer, IFCA (Santander)

Louvain, 05/2007

1. Introduction to SUSY
2. SUSY Lagrangians and the MSSM
3. “Simplified versions” and special sectors
4. SUSY Phenomenology

SUSY lectures (III): “Simplified Versions” and Special Sectors

Sven Heinemeyer, IFCA (Santander)

Louvain, 05/2007

1. Constrained Models:
 - CMSSM/mSUGRA
 - mGMSB
 - mAMSB
2. The MSSM Higgs Sector with real parameters
3. The MSSM Higgs Sector with complex parameters

1A. Gravity-mediated SUSY breaking

⇒ Quantum field theory of supergravity: graviton and gravitino

QFT with spin 2 and spin $\frac{3}{2}$ field is not renormalizable

⇒ cannot be extended to arbitrarily high energies

⇒ QFT of supergravity has to be interpreted as effective theory

contains non-renormalizable terms prop. to inverse powers of M_{Pl}

Best candidate for fundamental theory: string theory

SUSY breaking in hidden sector:

⇒ supergravity Lagrangian contains non-renormalizable terms that communicate between hidden and visible sector $\sim 1/M_{\text{Pl}}^n$

Dimensional analysis:

SUSY breaking in hidden sector by v.e.v. $\langle F \rangle$ ($\dim \langle F \rangle = \text{mass}^2$)
coupling $\sim 1/M_{\text{Pl}}$

require $m_{\text{soft}} \rightarrow 0$ for $\langle F \rangle \rightarrow 0$ (no SUSY breaking) and for
 $M_{\text{Pl}} \rightarrow \infty$ (vanishing gravitational interaction)

$$\Rightarrow m_{\text{soft}} \approx \frac{\langle F \rangle}{M_{\text{Pl}}}$$

Wanted: $m_{\text{soft}} \lesssim 1 \text{ TeV}$ (hierarchy problem)

$\Rightarrow \sqrt{\langle F \rangle} \approx 10^{11} \text{ GeV}$: scale of SUSY breaking in hidden sector

In general: $m_{\text{gravitino}} = m_{\frac{3}{2}} \approx \frac{\langle F \rangle}{M_{\text{Pl}}}$

$\Rightarrow m_{\frac{3}{2}} \approx m_{\text{soft}}$, gravitational interactions

\Rightarrow gravitino not important for collider phenomenology

Non-renormalizable terms in supergravity Lagrangian:

$$\begin{aligned}\mathcal{L}_{\text{NR}} = & -\frac{1}{M_{\text{Pl}}} F_X \sum_a \frac{1}{2} f_a \lambda^a \lambda^a + \text{h.c.} - \frac{1}{M_{\text{Pl}}^2} F_X F_X^* k_j^i \varphi_i \varphi^{*j} \\ & - \frac{1}{M_{\text{Pl}}} F_X \left(\frac{1}{6} y'^{ijk} \varphi_i \varphi_j \varphi_k + \frac{1}{2} \mu'^{ij} \varphi_i \varphi_j \right) + \text{h.c.}\end{aligned}$$

F_X : (auxiliary) field for a chiral supermultiplet X in the hidden sector

φ_i, λ^a : scalar and gaugino fields in the MSSM

If $\sqrt{\langle F_X \rangle} \sim 10^{10} - 10^{11}$ GeV

\Rightarrow soft SUSY-breaking terms of MSSM with $m_{\text{soft}} \approx 10^2 - 10^3$ GeV

Assumption of a “minimal” form of the supergravity Lagrangian

\Rightarrow soft-breaking terms which obey “universality” and “proportionality”

Results in exactly the known MSSM Lagrangian (??)

$$\begin{aligned}
 \mathcal{L}_{\text{soft}} = & -\frac{1}{2} \left(M_1 \tilde{B}\tilde{B} + M_2 \tilde{W}\tilde{W} + M_3 \tilde{g}\tilde{g} \right) + \text{h.c.} \\
 & - (m_{H_u}^2 + |\mu|^2) H_u^+ H_u - (m_{H_d}^2 + |\mu|^2) H_d^+ H_d - (b H_u H_d + \text{h.c.}) \\
 & - \left(\tilde{u}_R \mathbf{a}_u \tilde{Q} H_u - \tilde{d}_R \mathbf{a}_d \tilde{Q} H_d - \tilde{e}_R \mathbf{a}_e \tilde{L} H_d \right) + \text{h.c.} \\
 & - \tilde{Q}^+ \mathbf{m}_Q^2 \tilde{Q} - \tilde{L}^+ \mathbf{m}_L^2 \tilde{L} - \tilde{u}_R \mathbf{m}_u^2 \tilde{u}_R^* - \tilde{d}_R \mathbf{m}_d^2 \tilde{d}_R^* - \tilde{e}_R \mathbf{m}_e^2 \tilde{e}_R^*
 \end{aligned}$$

with 5 independent parameters at the GUT scale:

$$\begin{aligned}
 M_1 = M_2 = M_3 & = m_{1/2} \\
 m_{H_u}^2 = m_{H_d}^2 = \mathbf{m}_Q^2 = \mathbf{m}_L^2 = \mathbf{m}_u^2 = \mathbf{m}_d^2 = \mathbf{m}_e^2 & = m_0 \\
 \mathbf{a}_u = \mathbf{a}_d = \mathbf{a}_e & = A_0 \\
 & b \\
 & |\mu|^2
 \end{aligned}$$

Still to do: parameter(Q_{GUT}) \rightarrow parameter($Q_{\text{electroweak}}$)


Low-energy parameters (at the electroweak (EW) scale) via
 "Renormalization group equations" (RGEs)

[RGE: equations that connect parameters at different energy scales]

$\Rightarrow M_1, M_2, M_3, m_{H_u}^2, m_{H_d}^2, m_{\tilde{Q}}^2, m_{\tilde{L}}^2, m_{\tilde{u}}^2, m_{\tilde{d}}^2, m_{\tilde{e}}^2, a_u, a_d, a_e, b, |\mu|^2$ at the EW scale

Example:

$$M_0=300 \text{ GeV}, M_{1/2}=100 \text{ GeV}, A_0=0$$



Five new parameters, if possible phases are ignored:

$$m_0^2, m_{1/2}, A_0, b, \mu$$

Final "trick": require radiative electroweak symmetry breaking:

Require correct value of M_Z at the EW scale:

$$|\mu|^2 + m_{H_d}^2 = b \tan \beta - M_Z^2/2 \cos 2\beta$$

$$|\mu|^2 + m_{H_u}^2 = b \cot \beta + M_Z^2/2 \cos 2\beta$$

$\Rightarrow |\mu|, b$ given in terms of $\tan \beta, \text{sign } \mu$

\Rightarrow Scenario characterized by

$$m_0^2, m_{1/2}, A_0, \tan \beta, \text{sign } \mu$$

Usually called 'CMSSM' (constrained MSSM) or 'mSUGRA'

In agreement with all phenomenological constraints (see below)

Summary: “supergravity inspired scenario”, “mSUGRA”
characterized by five parameters:

$$m_0^2, m_{1/2}, A_0, \tan \beta, \text{sign } \mu$$

m_0 : universal scalar mass parameter

$m_{1/2}$: universal gaugino mass parameter

A_0 : universal trilinear coupling

$\tan \beta$: ratio of Higgs vacuum expectation values

$\text{sign}(\mu)$: sign of supersymmetric Higgs parameter

$m_0, m_{1/2}, A_0$: GUT scale parameters

⇒ particle spectra from renormalization group running to weak scale

Lightest SUSY particle (LSP) is usually lightest neutralino


gaugino masses run in same way as gauge couplings

⇒ gluino heavier than charginos, neutralinos

“Typical” mSUGRA scenario
 (SPS 1a benchmark scenario):

SPS home page:

www.ippp.dur.ac.uk/~georg/sps



1B. (minimal) gauge mediated SUSY breaking: mGMSB

New chiral supermultiplets, “messengers”, couple to SUSY breaking in hidden sector

Couple indirectly to MSSM fields via gauge interactions

⇒ mediation of SUSY breaking via electroweak and QCD gauge interactions

⇒ \approx flavor-diagonal

SUSY breaking already in messenger spectrum

⇒ masses of SUSY particles from loop diagrams with messenger particles, gauge-interaction strength

$$\Rightarrow m_{\text{soft}} \approx \frac{\alpha_i \langle F \rangle}{4\pi M_{\text{mess}}}, \quad M_{\text{mess}} \sim \sqrt{\langle F \rangle}$$


For $m_{\text{soft}} \lesssim 1 \text{ TeV} \Rightarrow \sqrt{\langle F \rangle} \approx 10^4\text{--}10^5 \text{ GeV}$

⇒ scale of SUSY breaking in hidden sector much lower than in SUGRA


Gravitino mass: $m_{\frac{3}{2}} \approx \frac{\langle F \rangle}{M_{\text{Pl}}} \approx 10^{-9} \text{ GeV}$

⇒ Gravitino is always the lightest SUSY particle (LSP)

Gaugino masses generated at one-loop order, $m_\lambda \approx \frac{\alpha_i}{4\pi}$



Scalar masses generated at two-loop order, $m_\varphi^2 \approx \left(\frac{\alpha_i}{4\pi}\right)^2$



⇒ Typical mass hierarchy in GMSB scenario between strongly interacting and weakly interacting particles $\sim \alpha_3/\alpha_2/\alpha_1$

GMSB scenario characterized by

$$M_{\text{mess}}, N_{\text{mess}}, \Lambda, \tan \beta, \text{sign}(\mu)$$

M_{mess} : messenger mass scale

N_{mess} : messenger index (number of messenger multiplets)


$\Lambda = \langle F \rangle / M_{\text{mess}}$: universal soft SUSY breaking mass scale
felt by low-energy sector

LSP is always the gravitino

next-to-lightest SUSY particle (NLSP): $\tilde{\chi}_1^0$ or $\tilde{\tau}_1$

can decay into LSP inside or outside the detector

GMSB scenario with $\tilde{\tau}$ NLSP
 (SPS 7 benchmark scenario):




1C. (minimal) anomaly mediated SUSY breaking: mAMSB

No MSSM in the “bulk”

Anomaly: $\langle F \rangle$ enters RGEs

$$m_{\tilde{f}_k}^2 \sim \frac{|\langle F \rangle|^2}{(16\pi^2)^2} g_k^4, \quad M_i \sim \frac{\langle F \rangle}{16\pi^2} g_i^2$$



mAMSB scenario characterized by

$$m_{3/2}, m_0, \tan \beta, \text{sign}(\mu)$$

$m_{3/2} = \langle F \rangle / M_{\text{Planck}}$: overall scale of SUSY particle masses


m_0 : phenomenological parameter: universal scalar mass term introduced in order to keep squares of slepton masses positive

AMSB scenario (SPS 9):

typical feature: very small neutralino–chargino mass difference


$$\Rightarrow \tilde{\chi}_1^\pm \rightarrow \tilde{\chi}_1^0 + \pi^\pm$$

with very soft pions



1.5 The Standard Model Higgs Sector


Current status of knowledge: the Standard Model (SM)



⇒ all particles experimentally seen

1.5 The Standard Model Higgs Sector

Current status of knowledge: the Standard Model (SM)



⇒ all particles experimentally seen

⇒ but one particle is missing . . .

Problem:

Gauge fields Z , W^+ , W^- are **massive**

explicit mass terms in the Lagrangian \Leftrightarrow breaking of gauge invariance

Solution: Higgs mechanism

scalar field postulated, mass terms from coupling to Higgs field

Higgs sector in the Standard Model:


Scalar SU(2) doublet: $\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}$

Higgs potential:

$$V(\phi) = \mu^2 |\Phi^\dagger \Phi| + \lambda |\Phi^\dagger \Phi|^2, \quad \lambda > 0$$

$\mu^2 < 0$: Spontaneous symmetry breaking

minimum of potential at $|\langle \Phi_0 \rangle| = \sqrt{\frac{-\mu^2}{2\lambda}} \equiv \frac{v}{\sqrt{2}}$



$$\Phi = \begin{pmatrix} 0 \\ v + H \end{pmatrix} \quad (\text{unitary gauge})$$

H : elementary scalar field, Higgs boson


Lagrange density:

$$\mathcal{L}_{\text{Higgs}} = (D_\mu \Phi)^\dagger (D^\mu \Phi) - V(\Phi)$$

Gauge invariant coupling to gauge fields

⇒ mass terms for gauge bosons and fermions

1.) $VV\Phi\Phi$ coupling:




⇒ VV mass terms

⇒ triple/quartic couplings to gauge bosons


$$g_2^2 v^2 / 2 \equiv M_W^2, \quad (g_1^2 + g_2^2) v^2 / 2 \equiv M_Z^2 \quad \Rightarrow \text{coupling} \propto \text{masses}$$

2.) fermion mass terms: Yukawa couplings



$$m_f = v g_f \Rightarrow \text{coupling} \propto \text{masses}$$

3.) mass of the Higgs boson: self coupling



$$\lambda = M_H^2/v$$

$$M_H = v\sqrt{\lambda} \quad \text{free parameter}$$

→ last unknown parameter of the SM

⇒ establish Higgs mechanism \equiv find the Higgs \oplus measure its couplings

Another effect of the Higgs field:

Scattering of longitudinal W bosons: $W_L W_L \rightarrow W_L W_L$

$$\mathcal{M}_V = \begin{array}{c} W \\ \diagup \\ \text{---} \\ \diagdown \\ W \end{array} \begin{array}{c} \text{---} \\ \diagup \\ \gamma, Z \\ \diagdown \\ \text{---} \end{array} \begin{array}{c} W \\ \diagdown \\ \text{---} \\ \diagup \\ W \end{array} + \begin{array}{c} \text{---} \\ \diagup \\ \gamma, Z \\ \diagdown \\ \text{---} \end{array} + \begin{array}{c} \text{---} \\ \diagup \\ \text{---} \\ \diagdown \\ \text{---} \end{array} = -g^2 \frac{E^2}{M_W^2} + \mathcal{O}(1) \quad \text{for } E \rightarrow \infty$$

\Rightarrow violation of unitarity

Contribution of a scalar particle with couplings prop. to the mass:

$$\mathcal{M}_S = \begin{array}{c} W \\ \diagup \\ \text{---} \\ \diagdown \\ W \end{array} \begin{array}{c} \text{---} \\ \diagup \\ H \\ \diagdown \\ \text{---} \end{array} \begin{array}{c} W \\ \diagdown \\ \text{---} \\ \diagup \\ W \end{array} + \begin{array}{c} \text{---} \\ \diagup \\ H \\ \diagdown \\ \text{---} \end{array} = g_{WWH}^2 \frac{E^2}{M_W^4} + \mathcal{O}(1) \quad \text{for } E \rightarrow \infty$$

$$\mathcal{M}_{\text{tot}} = \mathcal{M}_V + \mathcal{M}_S = \frac{E^2}{M_W^4} \left(g_{WWH}^2 - g^2 M_W^2 \right) + \dots$$


\Rightarrow compensation of terms with bad high-energy behavior for

$$g_{WWH} = g M_W$$

What else do we know about the Higgs boson?

SM at high energies

- upper limit on M_H :
 - dependence of coupling
 - λ_{HHHH} from energy scale Λ
 - ⇒ divergence: Landau pole
- lower limit on M_H :
 - stability of the vacuum :
 - $V(v) < V(0)$
 - [Coleman, Weinberg '73]
- combined ⇒




Λ : scale up to which the SM is valid


$$\Lambda = M_{\text{GUT}} \Rightarrow 130 \text{ GeV} \lesssim M_H \lesssim 180 \text{ GeV}$$

Indirect measurements via precision observables (POs):

Comparison of electro-weak precision observables with theory:



Test of theory at quantum level: Sensitivity to loop corrections



All parameters of the model enter
limits on M_H

Global fit to all SM data:


[LEPEWWG '07]

$$\Rightarrow M_H = 76^{+33}_{-24} \text{ GeV}$$

$$M_H < 144 \text{ GeV, 95\% C.L.}$$

Assumption for the fit:
SM incl. Higgs boson

\Rightarrow no confirmation of
Higgs mechanism



\Rightarrow Higgs boson seems to be light, $M_H \lesssim 150$ GeV

2. The MSSM Higgs Sector with real parameters

Two Higgs doublets:

$$H_1 = \begin{pmatrix} H_1^1 \\ H_1^2 \end{pmatrix} = \begin{pmatrix} v_1 + (\phi_1^0 - i\chi_1^0)/\sqrt{2} \\ -\phi_1^- \end{pmatrix}$$
$$H_2 = \begin{pmatrix} H_2^1 \\ H_2^2 \end{pmatrix} = \begin{pmatrix} \phi_2^+ \\ v_2 + (\phi_2^0 + i\chi_2^0)/\sqrt{2} \end{pmatrix}$$

→ Higgs potential (for the neutral Higgs bosons)

2. The MSSM Higgs Sector with real parameters

Two Higgs doublets:

$$H_1 = \begin{pmatrix} H_1^1 \\ H_1^2 \end{pmatrix} = \begin{pmatrix} v_1 + (\phi_1^0 - i\chi_1^0)/\sqrt{2} \\ -\phi_1^- \end{pmatrix}$$
$$H_2 = \begin{pmatrix} H_2^1 \\ H_2^2 \end{pmatrix} = \begin{pmatrix} \phi_2^+ \\ v_2 + (\phi_2^0 + i\chi_2^0)/\sqrt{2} \end{pmatrix}$$

Higgs potential:

$$V = m_1^2 H_1 \bar{H}_1 + m_2^2 H_2 \bar{H}_2 - m_{12}^2 (\epsilon_{ab} H_1^a H_2^b + \text{h.c.})$$
$$+ \underbrace{\frac{g'^2 + g^2}{8}}_{\text{gauge couplings, in contrast to SM}} (H_1 \bar{H}_1 - H_2 \bar{H}_2)^2 + \underbrace{\frac{g^2}{2}}_{\text{gauge couplings, in contrast to SM}} |H_1 \bar{H}_2|^2$$

Mixing of \mathcal{CP} -even, \mathcal{CP} -odd, charged fields:

$$\begin{pmatrix} H^0 \\ h^0 \end{pmatrix} = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} \phi_1^0 \\ \phi_2^0 \end{pmatrix}$$

$$\begin{pmatrix} G^0 \\ A^0 \end{pmatrix} = \begin{pmatrix} \cos \beta & \sin \beta \\ -\sin \beta & \cos \beta \end{pmatrix} \begin{pmatrix} \chi_1^0 \\ \chi_2^0 \end{pmatrix}, \quad \begin{pmatrix} G^\pm \\ H^\pm \end{pmatrix} = \begin{pmatrix} \cos \beta & \sin \beta \\ -\sin \beta & \cos \beta \end{pmatrix} \begin{pmatrix} \phi_1^\pm \\ \phi_2^\pm \end{pmatrix}$$

$$\tan(2\alpha) = \tan(2\beta) \frac{M_A^2 + M_Z^2}{M_A^2 - M_Z^2}$$

Three Goldstone bosons (as in SM): G^0, G^\pm

→ longitudinal components of W^\pm, Z

⇒ Five physical states: h^0, H^0, A^0, H^\pm

h, H : neutral, \mathcal{CP} -even, A^0 : neutral, \mathcal{CP} -odd, H^\pm : charged

Gauge-boson masses:

$$M_W^2 = \frac{1}{2}g'^2(v_1^2 + v_2^2), \quad M_Z^2 = \frac{1}{2}(g^2 + g'^2)(v_1^2 + v_2^2), \quad M_\gamma = 0$$

Problem:

MSSM contains term $\mu H_1 H_2$ in superpotential

μ : dimensionful parameter

For ew symmetry breaking required: $\mu \sim$ electroweak scale

But: no a priori reason for $\mu \neq 0$, $\mu \ll M_P$

(problem mainly in GMSB scenario, easier to overcome in mSUGRA)

Possible solution:

μ related to v.e.v. of additional field

\Rightarrow Introduction of extra singlet field S , v.e.v. $s \Rightarrow$ “NMSSM”

Superpotential: $\mathcal{V} = \lambda H_1 H_2 S + \frac{1}{3} \kappa S^3 + \dots$

Physical states in NMSSM Higgs-sector:

S_1, S_2, S_3 (CP-even), P_1, P_2 (CP-odd), H^\pm

Parameters in MSSM Higgs potential V (besides g, g'):

$$v_1, v_2, m_1, m_2, m_{12}$$

relation for $M_W^2, M_Z^2 \Rightarrow 1$ condition

minimization of V w.r.t. neutral Higgs fields $H_1^1, H_2^2 \Rightarrow 2$ conditions

\Rightarrow only two free parameters remain in V , conventionally chosen as

$$\tan \beta = \frac{v_2}{v_1}, \quad M_A^2 = -m_{12}^2(\tan \beta + \cot \beta)$$

$\Rightarrow m_h, m_H, \text{ mixing angle } \alpha, m_{H^\pm}$: no free parameters, can be predicted

In lowest order:

$$m_{H^\pm}^2 = M_A^2 + M_W^2$$

Predictions for m_h , m_H from diagonalization of tree-level mass matrix:

$\phi_1 - \phi_2$ basis:

$$M_{\text{Higgs}}^{2,\text{tree}} = \begin{pmatrix} m_{\phi_1}^2 & m_{\phi_1\phi_2}^2 \\ m_{\phi_1\phi_2}^2 & m_{\phi_2}^2 \end{pmatrix} =$$
$$\begin{pmatrix} M_A^2 \sin^2 \beta + M_Z^2 \cos^2 \beta & -(M_A^2 + M_Z^2) \sin \beta \cos \beta \\ -(M_A^2 + M_Z^2) \sin \beta \cos \beta & M_A^2 \cos^2 \beta + M_Z^2 \sin^2 \beta \end{pmatrix}$$

⇓ ← Diagonalization, α

$$\begin{pmatrix} m_H^{2,\text{tree}} & 0 \\ 0 & m_h^{2,\text{tree}} \end{pmatrix}$$

Tree-level result for m_h, m_H :

$$m_{H,h}^2 = \frac{1}{2} \left[M_A^2 + M_Z^2 \pm \sqrt{(M_A^2 + M_Z^2)^2 - 4M_Z^2 M_A^2 \cos^2 2\beta} \right]$$

$\Rightarrow m_h \leq M_Z$ at tree level

\Rightarrow Light Higgs boson h required in SUSY

Measurement of m_h , Higgs couplings

\Rightarrow test of the theory (more directly than in SM)

Higgs couplings, tree level:

$$g_{hVV} = \sin(\beta - \alpha) g_{HVV}^{\text{SM}}, \quad V = W^\pm, Z$$

$$g_{HVV} = \cos(\beta - \alpha) g_{HVV}^{\text{SM}}$$

$$g_{hAZ} = \cos(\beta - \alpha) \frac{g'}{2 \cos \theta_W}$$

$$g_{hb\bar{b}}, g_{h\tau^+\tau^-} = -\frac{\sin \alpha}{\cos \beta} g_{Hb\bar{b}, H\tau^+\tau^-}^{\text{SM}}$$

$$g_{ht\bar{t}} = \frac{\cos \alpha}{\sin \beta} g_{Ht\bar{t}}^{\text{SM}}$$

$$g_{Ab\bar{b}}, g_{A\tau^+\tau^-} = \gamma_5 \tan \beta g_{Hb\bar{b}}^{\text{SM}}$$

$\Rightarrow g_{hVV} \leq g_{HVV}^{\text{SM}}, \quad g_{hVV}, g_{HVV}, g_{hAZ}$ cannot all be small

$g_{hb\bar{b}}, g_{h\tau^+\tau^-}$: significant suppression or enhancement w.r.t. SM coupling possible

The decoupling limit:


For $M_A \gtrsim 150$ GeV:

The lightest MSSM Higgs is
SM-like

The heavy MSSM Higgses:

$$M_A \approx M_H \approx M_{H^\pm}$$

of course there are exceptions ...



Higgs mass bounds in SUSY theories

MSSM predicts upper bound on M_h :

tree-level bound: $m_h < M_Z$, excluded by LEP Higgs searches!

Large radiative corrections:

Yukawa couplings: $\frac{e m_t}{2M_W s_W}$, $\frac{e m_t^2}{M_W s_W}$, \dots

\Rightarrow Dominant one-loop corrections: $\Delta M_h^2 \sim G_\mu m_t^4 \log \left(\frac{m_{\tilde{t}_1} m_{\tilde{t}_2}}{m_t^2} \right)$

The MSSM Higgs sector is connected to all other sector via loop corrections (especially to the scalar top sector)

Present status of M_h prediction in the MSSM:

Complete one-loop and 'almost complete' two-loop result available

Upper bound on M_h in the MSSM:

“Unconstrained MSSM”:

M_A , $\tan \beta$, 5 parameters in \tilde{t} - \tilde{b} sector, μ , $m_{\tilde{g}}$, M_2

$$M_h \lesssim 135 \text{ GeV}$$

for $m_t = 170.9 \pm 1.8 \text{ GeV}$

(including theoretical uncertainties from unknown higher orders)

\Rightarrow observable at the LHC

Obtained with:

FeynHiggs

[S.H., W. Hollik, G. Weiglein '98, '00, '02]


[T. Hahn, S.H., W. Hollik, G. Weiglein '03 – '07]

www.feynhiggs.de

\rightarrow all Higgs masses, couplings, BRs (easy to link, easy to use :-)


Effects of the two-loop corrections to the lightest Higgs mass:

Example for one set of MSSM parameters



Effects of the two-loop corrections to the lightest Higgs mass:

Example for one set of MSSM parameters



Upper bound $M_h \lesssim 135$ GeV reduced by

$\approx 7, 12, 11$ GeV in **mSUGRA** , **GMSB** , **AMSB** scenarios

[*S. Ambrosanio, A. Dedes, S.H., S. Su, G. Weiglein'01*]

Upper bound on M_h in extensions of MSSM: $M_h \lesssim 200$ GeV
(no new gauge groups!)

[*G. Kane, C. Kolda, J. Wells '93*] [*J. Espinosa, M. Quirós '93, '98*]

\Rightarrow SUSY requires light Higgs boson:

definite and robust prediction of SUSY models

testable at next generation of colliders

Remaining theoretical uncertainties in prediction for M_h in the MSSM:

[G. Degrandi, S.H., W. Hollik, P. Slavich, G. Weiglein '02]

- From unknown higher-order corrections:
 $\Rightarrow \Delta M_h \approx 3 \text{ GeV}$

- From uncertainties in input parameters

$m_t, \dots, M_A, \tan \beta, m_{\tilde{t}_1}, m_{\tilde{t}_2}, \theta_{\tilde{t}}, m_{\tilde{g}}, \dots$

$\Delta m_t \approx 2 \text{ GeV} \Rightarrow \Delta M_h \approx 2 \text{ GeV}$

Higgs couplings, production cross sections


Also affected by large SUSY loop corrections

FD two-loop corrections implemented in results for $h \rightarrow f\bar{f}, e^+e^- \rightarrow hZ, hA$

[S.H., W. Hollik, G. Weiglein '00]

[S.H., W. Hollik, J. Rosiek, G. Weiglein '01]

$hf\bar{f}$ coupling:



$$A(h \rightarrow f\bar{f}) = \sqrt{Z_h} \left(\Gamma_h - \frac{\hat{\Sigma}_{hH}(M_h^2)}{M_h^2 - m_H^2 + \hat{\Sigma}_{HH}(M_h^2)} \Gamma_H \right)$$

\Rightarrow Effective $hf\bar{f}$ coupling can vanish for large $\hat{\Sigma}_{hH}$

Glauino vertex corrections to $h \rightarrow q\bar{q}$:

\Rightarrow ratio $\Gamma(h \rightarrow \tau^+\tau^-)/\Gamma(h \rightarrow b\bar{b})$ can significantly differ from SM value for large $\tan\beta$


Effective $hf\bar{f}$ coupling can go to zero for large $\hat{\Sigma}_{hH}$

⇒ “Pathological regions”

[W. Loinaz, J. Wells '98] [M. Carena, S. Mrenna, C. Wagner '99]

⇒ Suppression of $BR(h \rightarrow b\bar{b})$,
 $BR(h \rightarrow \tau\tau)$, ...

[S.H., W. Hollik, G. Weiglein '00]



3. The Higgs sector in the MSSM with complex parameters (cMSSM)

Higgs potential of the cMSSM contains two Higgs doublets:

$$H_1 = \begin{pmatrix} H_1^1 \\ H_1^2 \end{pmatrix} = \begin{pmatrix} v_1 + (\phi_1 + i\chi_1)/\sqrt{2} \\ \phi_1^- \end{pmatrix}$$
$$H_2 = \begin{pmatrix} H_2^1 \\ H_2^2 \end{pmatrix} = e^{i\xi} \begin{pmatrix} \phi_2^+ \\ v_2 + (\phi_2 + i\chi_2)/\sqrt{2} \end{pmatrix}$$

$$V = m_1^2 H_1 \bar{H}_1 + m_2^2 H_2 \bar{H}_2 - (m_{12}^2 \epsilon_{ab} H_1^a H_2^b + \text{h.c.})$$
$$+ \underbrace{\frac{g'^2 + g^2}{8}}_{\text{gauge couplings, in contrast to SM}} (H_1 \bar{H}_1 - H_2 \bar{H}_2)^2 + \underbrace{\frac{g^2}{2}}_{\text{gauge couplings, in contrast to SM}} |H_1 \bar{H}_2|^2$$

Five physical states: h^0, H^0, A^0, H^\pm (no CPV at tree-level)

2 CP-violating phases: $\xi, \arg(m_{12}) \Rightarrow$ can be set/rotated to zero

Input parameters: $\tan \beta = \frac{v_2}{v_1}, M_A$ or M_{H^\pm}

Effects of complex parameters in the Higgs sector:

Complex parameters enter via loop corrections:

- μ : Higgsino mass parameter
- $A_{t,b,\tau}$: trilinear couplings $\Rightarrow X_{t,b,\tau} = A_{t,b,\tau} - \mu^* \{\cot \beta, \tan \beta\}$ complex
- $M_{1,2}$: gaugino mass parameter (one phase can be eliminated)
- M_3 : gluino mass parameter

\Rightarrow can induce \mathcal{CP} -violating effects

Result:

$$(A, H, h) \rightarrow (h_3, h_2, h_1)$$

with

$$M_{h_3} > M_{h_2} > M_{h_1}$$

Inclusion of higher-order corrections:

(→ Feynman-diagrammatic approach)

Propagator / mass matrix with higher-order corrections:

$$\begin{pmatrix} q^2 - M_A^2 + \hat{\Sigma}_{AA}(q^2) & \hat{\Sigma}_{AH}(q^2) & \hat{\Sigma}_{Ah}(q^2) \\ \hat{\Sigma}_{HA}(q^2) & q^2 - m_H^2 + \hat{\Sigma}_{HH}(q^2) & \hat{\Sigma}_{Hh}(q^2) \\ \hat{\Sigma}_{hA}(q^2) & \hat{\Sigma}_{hH}(q^2) & q^2 - m_h^2 + \hat{\Sigma}_{hh}(q^2) \end{pmatrix}$$

$\hat{\Sigma}_{ij}(q^2)$ ($i, j = h, H, A$) : renormalized Higgs self-energies

$\hat{\Sigma}_{Ah}, \hat{\Sigma}_{AH} \neq 0 \Rightarrow \mathcal{CPV}$, \mathcal{CP} -even and \mathcal{CP} -odd fields can mix

Results available for $\hat{\Sigma}_{ij}$:


– full 1-loop evaluation: dependence on all possible phases included

– New: $\mathcal{O}(\alpha_t \alpha_s)$ corrections in the FD approach

(rMSSM: difference between FD and RGiEP approach $\mathcal{O}(\text{few GeV})$)

M_{h_1} as a function of ϕ_{A_t} :

[S.H., W. Hollik, H. Rzehak, G. Weiglein '05]



$$M_{\text{SUSY}} = 1000 \text{ GeV}$$

$$|A_t| = 2000 \text{ GeV}$$

$$\tan \beta = 10$$


$$M_{H^\pm} = 150 \text{ GeV}$$

OS renormalization

\Rightarrow modified dependence
on ϕ_{A_t} at the 2-loop level

M_{h_1} as a function of $\phi_{\tilde{g}}$:

[S.H., W. Hollik, H. Rzehak, G. Weiglein '05]



$$M_{\text{SUSY}} = 500 \text{ GeV}$$

$$A_t = 1000 \text{ GeV}$$

$$\tan \beta = 10$$

$$M_{H^\pm} = 500 \text{ GeV}$$

OS renormalization

\Rightarrow threshold at $m_{\tilde{g}} = m_{\tilde{t}} + m_t$

\Rightarrow large effects around threshold

\Rightarrow phase dependence has to be taken into account

4. The code FeynHiggs 2.5

Latest version: FeynHiggs 2.5 (11/06)

real MSSM:

contains all available higher-order corrections
to Higgs boson masses and couplings

FeynHiggs contains

- full 1 loop calculations
- all available 2 loop calculations (leading and subleading)
- very leading 3 loop contributions

complex MSSM:

contains nearly all available results
(we are (even currently) working on the rest)

www.feynhiggs.de

Included in FeynHiggs 2.5 (I):

Evaluation of all Higgs boson masses and mixing angles

- $M_{h_1}, M_{h_2}, M_{h_3}, M_{H^\pm}$, α_{eff} , \mathbf{Z}_{ij} , \mathbf{U}_{ij} , ...

Evaluation of all neutral Higgs boson decay channels \Leftarrow with \mathbf{Z}

- total decay width Γ_{tot}
- $\text{BR}(h_i \rightarrow f\bar{f})$: decay to SM fermions
- $\text{BR}(h_i \rightarrow \gamma\gamma, ZZ^{(*)}, WW^{(*)}, gg)$: decay to SM gauge bosons
- $\text{BR}(h_i \rightarrow h_1 Z^{(*)}, h_1 h_1)$: decay to gauge and Higgs bosons
- $\text{BR}(h_i \rightarrow \tilde{f}_i \tilde{f}_j)$: decay to sfermions
- $\text{BR}(h_i \rightarrow \tilde{\chi}_i^\pm \tilde{\chi}_j^\pm, \tilde{\chi}_i^0 \tilde{\chi}_j^0)$: decay to charginos, neutralinos

Evaluation for the SM Higgs (same masses as the three MSSM Higgses)

- total decay width $\Gamma_{\text{tot}}^{\text{SM}}$
- $\text{BR}(h_i^{\text{SM}} \rightarrow f\bar{f})$: decay to SM fermions
- $\text{BR}(h_i^{\text{SM}} \rightarrow \gamma\gamma, ZZ^{(*)}, WW^{(*)}, gg)$: decay to SM gauge bosons

Included in FeynHiggs 2.5 (II):

Evaluation of all neutral Higgs boson production cross sections at Tevatron/LHC \Leftarrow with \mathbf{Z}

SM: most up-to-date, MSSM: additional effective couplings

- $gg \rightarrow h_i$: gluon fusion
- $WW \rightarrow h_i, ZZ \rightarrow h_i$: gauge boson fusion
- $W \rightarrow Wh_i, Z \rightarrow Zh_i$: Higgs strahlung
- $b\bar{b} \rightarrow b\bar{b}h_i$: Yukawa process
- $b\bar{b} \rightarrow b\bar{b}h_i, h_i \rightarrow b\bar{b}$, one b tagged
- $t\bar{t} \rightarrow t\bar{t}h_i$: Yukawa process
- $\tilde{t}_1\tilde{t}_1 \rightarrow \tilde{t}_1\tilde{t}_1h_i$: SUSY-Yukawa process

Evaluation for the SM Higgs (same masses as the three MSSM Higgses)

- all SM-like channels as above

Included in FeynHiggs 2.5 (III):

Evaluation of all charged Higgs boson decay channels (rMSSM/cMSSM)

- total decay width Γ_{tot}
- $\text{BR}(H^+ \rightarrow f\bar{f}')$: decay to SM fermions
- $\text{BR}(H^+ \rightarrow h_i W^+)$: decay to gauge and Higgs bosons
- $\text{BR}(H^+ \rightarrow \tilde{f}_i \tilde{f}'_j)$: decay to sfermions
- $\text{BR}(H^+ \rightarrow \tilde{\chi}_i^0 \tilde{\chi}_j^+)$: decay to charginos and neutralinos

Evaluation of additional couplings: \Leftarrow with **U**

- $g(V \rightarrow V h_i, h_i h_j)$: coupling of gauge and Higgs bosons
- $g(h_i h_j h_k)$: all Higgs self couplings (including charged Higgs)
- $\sigma(\gamma\gamma \rightarrow h_i)$: Higgs production XS at a γC

Included in FeynHiggs 2.5 (IV):

Evaluation of theory error on masses and mixing

→ estimate of uncertainty in $M_{h_i}, \mathbf{U}_{ij}, \mathbf{Z}_{ij}$ from unknown higher-order corr.

Evaluation of masses, mixing and decay in the NMFV MSSM

NMFV: Non Minimal Flavor Violation [Hahn, S.H., Hollik, Merz, Peñaranda '04-'06]
⇒ Connection to Flavor physics

Evaluation of additional constraints (rMSSM/cMSSM)

- ρ -parameter: $\Delta\rho^{\text{SUSY}}$ at $\mathcal{O}(\alpha), \mathcal{O}(\alpha\alpha_s), \dots$, including NMFV effects
⇒ $M_W, \sin^2\theta_{\text{eff}}$ via SM formula + $\Delta\rho^{\text{SUSY}}$, including NMFV effects
- anomalous magnetic moment of the μ : $(g - 2)_\mu$
- $\text{BR}(b \rightarrow s\gamma)$, including NMFV effects [T. Hahn, W. Hollik, J. Illana, S. Peñaranda '06]
- LEP Higgs constraints [LEP Higgs WG '06]
- EDMs of electron, neutron, Hg, ...

Planned:

- ILC production cross sections

How to install FeynHiggs 2.5

1. Go to www.feynhiggs.de
2. Download the latest version
3. type `./configure, make, make install`
⇒ library `libFH.a` is created
4. 4 possible ways to use *FeynHiggs*:
 - A) Command-line mode
 - B) called from a Fortran/C++ code
 - C) called within *Mathematica*
 - D) *WWW* modeprocessing of *Les Houches Accord* data possible
5. Detailed `instructions` and `help` are provided in the `man pages`

How to run FeynHiggs 2.5

A) Command-line mode

Input File

```
MT      172.7
MB       4.7
MW      80.4
MZ      91.1
MSusy   975
MAO     200
Abs(M_2) 332
Abs(MUE) 980
TB       50
Abs(At) -300
Abs(Ab)  1500
Abs(M_3) 975
```

Command

FeynHiggs file flags

Screen Output

```
----- HIGGS MASSES -----
| Mh0    = 116.022817
| MHH    = 199.943497
| MA0    = 200.000000
| MHp    = 216.973920
| SAeff  = -0.02685112
| UHiggs = 0.99999346 -0.00361740 0.00000000 \
|        0.00361740 0.99999346 0.00000000 \
|        0.00000000 0.00000000 1.00000000
----- ESTIMATED UNCERTAINTIES -----
| DeltaMh0 = 1.591957
| DeltaMHH = 0.004428
| DeltaMA0 = 0.000000
| DeltaMHp = 0.152519
| ...
```

- Loops over parameter values possible (parameter scans).
- Mask off details with `FeynHiggs file flags | grep -v %`
- `table` utility converts to machine-readable format, e.g.
`FeynHiggs file flags | table TB Mh0 > outfile`

Input File

```
BLOCK MODSEL
  1  1
BLOCK MINPAR
  1  0.10000E+03 # m0
  2  0.25000E+03 # m12
  3  0.10000E+02 # tanb
  4  0.10000E+01 # sgn mu
  5 -0.10000E+03 # A
BLOCK SMINPUTS
  4  0.91187E+02 # MZ
  5  0.42500E+01 # mb(mb)
  6  0.17500E+03 # t
...
```

Command
FeynHiggs file flags

file.fh

```
BLOCK MASS
  25  1.12697840E+02 # Mh0
  35  4.00145460E+02 # MHH
  36  3.99769788E+02 # MA0
  37  4.08050556E+02 # MHp
  ...
BLOCK ALPHA
      -1.10658125E-01 # Alpha
  ...
```

- { Uses / was developed into } the SLHA I/O Library. [*T. Hahn '04*]
- SLHA can also be used in Library Mode with `FHSetSLHA`.
- *FeynHiggs* tries to read each file in SLHA format first. If that fails, fallback to native format.

B) Called from a Fortran/C++ code

Link *FeynHiggs* as a subroutine \Rightarrow link `libFH.a`

`call FHSetFlags(...)` :

→ specification of accuracy etc.

`call FHSetPara(...)` :

→ specify input parameters

`call FHGetPara(...)` :

→ obtain derived parameters

`call FHHiggsCorr(...)` :

→ obtain Higgs boson masses and mixings

`call FHUncertainties(...)` :

→ obtain theory error on Higgs boson masses and mixings from unknown higher-order corrections

`call FHCouplings(...)` :

→ obtain decay widths, BRs, XSs, etc.

C) Called within Mathematica

- install the `math link` to `MFeynHiggs` , e.g.:

```
Install[,'MFeynHiggs']
```

- `FHSetFlags[...]` :
→ specification of accuracy etc.

`FHSetPara[...]` :
→ specify input parameters

`FHGetPara[]` :
→ obtain derived parameters

`FHHiggsCorr[]` :
→ obtain Higgs boson masses and mixings

`FHUncertainties[]` :
→ obtain theory error on Higgs boson masses and mixings from
unknown higher-order corrections

`FHCouplings[]` :
→ obtain decay widths, BRs etc.

D) WWW mode

1. The **FeynHiggs User Control Center** is available at
www.feynhiggs.de/fhucc
2. Enter you parameters on-line in the web page
3. Obtain your **results with a mouse click**

⇒ for single points and checks of your downloaded version of FeynHiggs
⇒ always the latest version

⇒ online presentation

Also man pages are available on-line

D) WWW mode

1. The FeynHiggs User Control Center is available at

2. <http://www.feynhiggs.de/cgi-bin/ft> /fhucc

3. **The FeynHiggs User Control Center**

⇒ **Flags**

⇒ Scope of the 1-loop part: full MSSM

1-loop field renormalization: MSbar

FeynHiggs

Also man pages are available on-line