




# High-spin scattering at 2PM

Kays Haddad  
November 29, 2022

2203.06197 and 2205.02809 with Rafael Aoude & Andreas Helset



[LIGO Collaboration, '16]



weak gravity

$$\frac{GM}{R} \ll 1$$

non-relativistic

$$v \ll 1$$

weak gravity




post-Minkowskian (PM)  
expansion

weak gravity  
non-relativistic  
virial theorem




post-Newtonian (PN)  
expansion

		0PN	1PN	2PN	3PN	
0PM	1	$v^2$	$v^4$	$v^6$	$v^8$	...
1PM		$G$	$Gv^2$	$Gv^4$	$Gv^6$	...
2PM			$G^2$	$G^2v^2$	$G^2v^4$	...
3PM				$G^3$	$G^3v^2$	...
4PM					$G^4$	...




$\sim$



# Amplitudes and classical physics

conventional wisdom: trees are classical...



... and loops are quantum  $\sim \hbar^{L-1}$  [Itzykson, Zuber, '80].


counterexample:

$$\mathcal{A} = \underbrace{\frac{e^2 p_1 \cdot p_2}{\hbar m_1 m_2 q^2}}_{[E]^{-2}} + \underbrace{\frac{e^4}{16 m_1 m_2 \sqrt{-q^2}} (m_1 + m_2)}_{[L]^2}$$

to account for classical physics [Holstein, Donoghue, '04; Kosower, Maybee, O'Connell, '18],


$$e \rightarrow e/\sqrt{\hbar}, \quad G \rightarrow G/\hbar, \quad q^\mu \rightarrow \hbar \bar{q}^\mu$$

→ 
$$\mathcal{A} = \frac{e^2 p_1 \cdot p_2}{\hbar^3 m_1 m_2 q^2} + \frac{e^4}{16 \hbar^3 m_1 m_2 \sqrt{-q^2}} (m_1 + m_2)$$




## why amplitudes?


double copy:




unitarity/factorization:







# How do we model Kerr scattering?





$$\text{observable} \sim f_0 + f_1 \mathfrak{a} + f_2 \mathfrak{a}^2 + \dots$$



$$\text{observable} \sim g_0 + g_1 \frac{\langle S \rangle}{m} + g_2 \frac{\langle S^2 \rangle}{m^2} + \dots + g_{2s} \frac{\langle S^{2s} \rangle}{m^{2s}}$$

$$\text{classical limit, } \hbar \rightarrow 0, s \rightarrow \infty : \lim_{\hbar \rightarrow 0} \langle S^n \rangle = (m\mathfrak{a})^n, \quad \frac{d}{ds} \left( \lim_{\hbar \rightarrow 0} g_i \right) = 0, \quad \lim_{\hbar \rightarrow 0} g_i = f_i$$



$n$  gravitons

$n-1$  loops

$\mathcal{O}(G^n v^\infty)$

$n$ PM order


The diagram illustrates the expansion of a two-loop bubble diagram. On the left, two dashed circles (bubbles) are connected by a horizontal wavy line. Each bubble has two external red lines: one entering from the bottom and one exiting from the top. An arrow labeled  $q^\mu$  points from the left bubble to the right bubble. This is set equal to the product of two one-loop bubble diagrams, each with a single external red line, also connected by a wavy line and labeled with  $q^\mu$ . The final term is  $+ \mathcal{O}((q^2)^0)$ .

“minimal coupling” amplitude:  
 [Arkani-Hamed, Huang, Huang, ‘17]

$$\mathcal{M}_3^{(s)} = \frac{\langle \mathbf{11}' \rangle^{\odot 2s}}{m^{2s}} \mathcal{M}_3^{(0)}$$

[Guevara, Ochirov, Vines, ‘18] based on [Vines, ‘17]  
 [Chung, Huang, Kim, Lee, ‘18] based on [Levi, Steinhoff, ‘15]

$$\lim_{s \rightarrow \infty} \mathcal{M}_3^{(s)} = \mathcal{M}_3^{\text{Kerr}} = e^{\pm q \cdot \mathfrak{a}} \mathcal{M}_3^{\text{Schw}}$$




$$\mathcal{M}_{4,-+}^s = \frac{y^4}{s_{34}t_{13}t_{14}} \left( \frac{\langle 31 \rangle [42] - \langle 32 \rangle [41]}{y} \right)^{\odot 2s} \quad [\text{Arkani-Hamed, Huang, Huang, '17}]$$

$$\lim_{\substack{\hbar \rightarrow 0 \\ s \rightarrow \infty}} \mathcal{M}_{4,-+}^s = \frac{y^4}{s_{34}t_{13}t_{14}} \exp \left[ \left( q_4 - q_3 + \frac{t_{14} - t_{13}}{y} w \right) \cdot \mathbf{a} \right] \quad [\text{Aoude, KH, Helset, '20, '22}]$$

(underlying formalism: on-shell heavy spinors)

see also [Guevara, Ochirov, Vines, '18]

$$w^\mu \equiv \frac{1}{2} [4 | \bar{\sigma}^\mu | 3 \rangle, \quad y \equiv 2p_1 \cdot w$$



[Saketh, Vines, '22]  
[Fabian Bautista's thesis, '22]

$$\mathcal{M}_{4,-+}^\infty \supset \frac{(t_{14} - t_{13})^n}{s_{34} t_{13} t_{14}} \frac{(w \cdot \mathfrak{a})^n}{y^{n-4}}, \quad \mathcal{O}(\mathfrak{a}^{n > 4})$$


# A viable Compton amplitude

1. no spurious poles
2. factorizes onto three-point Kerr

see also [Chung, Huang, Kim, Lee, '18; Falkowski, Machado, '20; Chiodaroli, Johansson, Pichini, '21]



# QED




$$\begin{aligned}\mathcal{A}_{4,-+}^{\infty} &= \frac{y^2}{t_{13}t_{14}} \exp \left[ \left( q_4 - q_3 + \frac{t_{14} - t_{13}}{y} w \right) \cdot \mathbf{a} \right] \\ &= e^{(q_4 - q_3) \cdot \mathbf{a}} \sum_{n=0}^{\infty} \frac{1}{n!} I_n\end{aligned}$$

$$I_n \equiv \frac{y^2}{t_{13}t_{14}} \left( \frac{t_{14} - t_{13}}{y} w \cdot \mathbf{a} \right)^n$$


$$I_{n \geq 2} = -4 \left( \frac{t_{14} - t_{13}}{y} \right)^{n-2} (w \cdot \mathbf{a})^n$$

QED



$$\mathcal{A}_{4,-+}^{\infty} = e^{(q_4 - q_3) \cdot \mathbf{a}} \sum_{n=0}^2 \frac{1}{n!} I_n$$

# Gravity



$$\mathcal{M}_{4,-+}^{\infty} = e^{(q_4 - q_3) \cdot \mathbf{a}} \sum_{n=0}^{\infty} \frac{1}{n!} K_n$$

$$K_n = \frac{y^4}{s_{34} t_{13} t_{14}} \left( \frac{t_{14} - t_{13}}{y} w \cdot \mathbf{a} \right)^n$$

$$K_{n>4} = -\frac{4}{s_{34}} \frac{(t_{14} - t_{13})^{n-2}}{y^{n-4}} (w \cdot \mathbf{a})^n$$

# Gravity

vanishing Gram determinant:

$$K_{n \geq 5} = 16m^2 \frac{(t_{14} - t_{13})^{n-4}}{y^{n-4}} (w \cdot \mathfrak{a})^n + 2\mathfrak{s}_1 K_{n-1} - \mathfrak{s}_2 K_{n-2}$$


$$\begin{aligned}\mathfrak{s}_1 &\equiv (q_3 - q_4) \cdot \mathfrak{a}, \\ \mathfrak{s}_2 &\equiv -4(q_3 \cdot \mathfrak{a})(q_4 \cdot \mathfrak{a}) + s_{34}\mathfrak{a}^2\end{aligned}$$

solution:

$$\bar{K}_n \equiv \begin{cases} K_n, & n \leq 4, \\ K_4 L_{n-4} - K_3 \mathfrak{s}_2 L_{n-5}, & n > 4. \end{cases}$$

$$L_m \equiv \sum_{j=0}^{\lfloor m/2 \rfloor} \binom{m+1}{2j+1} \mathfrak{s}_1^{m-2j} (\mathfrak{s}_1^2 - \mathfrak{s}_2)^j$$

# Gravity



$$\mathcal{M}_{4,-+}^{\infty} = e^{(q_4 - q_3) \cdot \mathbf{a}} \sum_{n=0}^{\infty} \frac{1}{n!} \bar{K}_n$$

$$\mathcal{M}_{4,-+}^{\text{Kerr}} = \mathcal{M}_{4,-+}^{\infty} + m^2 \mathcal{C} \text{ where } \text{Res}_{t_{13}=0} \mathcal{C} = \text{Res}_{t_{14}=0} \mathcal{C} = \text{Res}_{s_{34}=0} \mathcal{C} = 0$$

# Contact terms\*

\*not necessarily describing Kerr

$$\mathcal{M}_{4,-+}^{\infty} + m^2 \mathcal{C}$$

decompose contact term into Wilson coefficients and functions of momenta and spin vector:

$$\mathcal{C} = \sum_{j,k} c_{j,k} \mathcal{C}_j(\mathbf{a}^k)$$

three relevant scales:  $c_{j,k} \sim G^{n_1} m^{n_2} \hbar^{n_3}$

$$\left. \begin{array}{l} \mathcal{M}_{4,-+}^{\infty} \sim \mathcal{O}(\hbar^0) \\ [\mathcal{M}_{4,-+}^{\infty}] = [m]^2 \end{array} \right\} \longrightarrow c_{j,k} \sim \left( \frac{Gm}{\hbar} \right)^n$$

$$\mathcal{C} = \sum_{j,k} c_{j,k} \mathcal{C}_j(\mathbf{a}^k) \quad c_{j,k} \sim \left( \frac{Gm}{\hbar} \right)^n$$

relevant at  $\mathcal{O}(G^2) \Rightarrow n = 0$

$$\longrightarrow \left\{ \begin{array}{l} c_{j,k} \mathcal{C}_j(\mathbf{a}^k) \Big|_{k \leq 3} = \mathcal{O}(\hbar) \\ 1 \leq j \leq \frac{1}{24}(2k^3 + 9k^2 - 74k) + \begin{cases} 4, & k \text{ even} \geq 4, \\ \frac{87}{24}, & k \text{ odd} \geq 5. \end{cases} \end{array} \right.$$



Kerr Compton amplitude for  $\mathcal{O}(\mathfrak{a}^{n \leq 4})$  invariant under shift (see also [Bern, Kosmopoulos, Luna, Roiban, Teng, '22])

$$\mathfrak{a}^\mu \rightarrow \mathfrak{a}^\mu + \xi(q_3^\mu + q_4^\mu)$$

require same symmetry of contact terms:

$$\mathcal{C} \equiv (w \cdot \mathfrak{a})^4 \sum_{n=0}^{\infty} \sum_{j=0}^{\infty} d_{n,j} \mathfrak{s}_1^n (\mathfrak{s}_1^2 - \mathfrak{s}_2)^j$$

only  $\lfloor k/2 \rfloor - 1$  terms at  $\mathcal{O}(\mathfrak{a}^{k \geq 4})$ .

# High-spin scattering at 2PM

spinning object  $\times$  Schwarzschild

$$\begin{array}{c} \mathcal{M}_{3,\pm}^{\text{Kerr}} \\ \mathcal{M}_{3,\mp}^{\text{Kerr}} \end{array} \begin{array}{c} \uparrow \\ \uparrow \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} \mathcal{M}_{4,\mp\pm}^{\infty} + \begin{array}{c} \uparrow \\ \uparrow \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} \begin{array}{c} \text{---} \\ \text{---} \end{array} \begin{array}{c} \uparrow \\ \uparrow \end{array}$$

no  $\mathcal{M}_{4,\pm\pm}^{\infty}$  if shift symmetry extends to 2PM

$$\mathcal{M}_{2\text{PM}} = G^2 m_1^2 m_2^2 \frac{\pi^2}{\sqrt{-q^2}} \sum_{n=0}^{\infty} \sum_{k=0}^n \left( M_k^{(2n)} + i\omega \mathcal{E}_1 M_k^{(2n+1)} \right) Q^{n-k} V^k$$

$$\omega \equiv v_1 \cdot v_2$$

$$\mathcal{E}_1 \equiv \epsilon^{\mu\nu\alpha\beta} v_{1\mu} v_{2\nu} q_\alpha \mathbf{a}_{1\beta}$$

$$Q \equiv (q \cdot \mathbf{a}_1)^2 - q^2 \mathbf{a}_1^2$$

$$V \equiv q^2 (v_2 \cdot \mathbf{a}_1)^2$$

fifth order in spin:

$$M_0^{(5)} = \frac{3m_2(4\omega^2 - 1) + 2m_1(13\omega^2 - 7)}{48(\omega^2 - 1)},$$

$$M_1^{(5)} = -\frac{m_2(5\omega^2 + 1) + 8m_1(2\omega^2 - 1)}{8(\omega^2 - 1)^2},$$

$$M_2^{(5)} = \frac{m_2(-7\omega^4 + 34\omega^2 - 3) + 32m_1(2\omega^2 - 1)}{48(\omega^2 - 1)^3}.$$

agrees with [Bern, Kosmopoulos, Luna, Roiban, Teng, '22]

seventh order in spin:

$$M_0^{(7)} = \frac{6m_2(8\omega^2 - 1) + 7m_1(17\omega^2 - 9)}{8064(\omega^2 - 1)},$$

$$M_1^{(7)} = -\frac{m_2(5\omega^2 + 1) + 8m_1(2\omega^2 - 1)}{192(\omega^2 - 1)^2},$$

$$M_2^{(7)} = \frac{m_2(-7\omega^4 + 34\omega^2 - 3) + 32m_1(2\omega^2 - 1)}{576(\omega^2 - 1)^3},$$

$$M_3^{(7)} = -\frac{m_2(9\omega^6 - 41\omega^4 + 95\omega^2 - 15) + 64m_1(2\omega^2 - 1)}{2880(\omega^2 - 1)^4}.$$

fourth order in spin:

$$M_0^{(4)} = \frac{m_1(239\omega^4 - 250\omega^2 + 35) + 3m_2 [8\omega^2(5\omega^2 - 4) + 3d_{0,0}(\omega^2 - 1)]}{96(\omega^2 - 1)},$$

$$M_1^{(4)} = -\frac{2m_1(193\omega^4 - 194\omega^2 + 25) - 3m_2 [8(-5\omega^4 + \omega^2 + 2) + 15d_{0,0}(\omega^2 - 1)^2]}{48(\omega^2 - 1)^2},$$

$$M_2^{(4)} = \frac{64m_1(8\omega^4 - 8\omega^2 + 1) + m_2 [8(15\omega^4 + 6\omega^2 - 13) + 105d_{0,0}(\omega^2 - 1)^3]}{96(\omega^2 - 1)^3}.$$

Kerr has  $d_{0,0} = 0$  which improves  $\lim_{\omega \rightarrow \infty} M_2^{(4)}$ ; agrees with [Chen, Chung, Huang, Kim, '21].


requiring best  $\lim_{\omega \rightarrow \infty} M_i^{(2n)}$  sets  $d_{2k,j} = -\frac{16(k-j)(2k+1)}{(2j+2k+4)!}$ .

$$\mathcal{M}_{2\text{PM}} = \frac{2G^2\pi^2 m_1^2 m_2^2}{\sqrt{-q^2}} (\mathcal{M}_{2\text{PM}}^{\text{even}} + i\omega \mathcal{E}_1 \mathcal{M}_{2\text{PM}}^{\text{odd}})$$

$$\begin{aligned} \mathcal{M}_{2\text{PM}}^{\text{even}} = & m_1 \left[ 3(5\omega^2 - 1)\mathcal{F}_0 + \frac{1}{4}(\omega^2 - 1)\mathcal{F}_2 Q + \frac{8\omega^4 - 8\omega^2 + 1}{\omega^2 - 1}\mathcal{F}_1 Q \right. \\ & \left. - \frac{1}{2}\mathcal{F}_2 V + \sum_{k=1}^{\infty} \frac{(8\omega^4 - 8\omega^2 + 1)}{(\omega^2 - 1)^{k+1}} \frac{(-1)^k 2\Gamma[k]}{\Gamma[2k]} \mathcal{F}_{k-1} V^k \right] \\ & - m_2 \left[ -3(5\omega^2 - 1)\sqrt{\pi}\mathcal{F}_{-1/2} - \frac{3\sqrt{\pi}}{4}\mathcal{F}_{1/2} Q - \frac{1}{\omega^2 - 1}\mathcal{F}_1 Q + \frac{15\sqrt{\pi}}{4}\mathcal{F}_{1/2} V \right. \\ & \left. + 6\sqrt{\pi} \sum_{k=1}^{\infty} \frac{\omega^{2k}}{(\omega^2 - 1)^{k+1}} \frac{(-1)^k \mathcal{F}_{k-1} V^k}{\Gamma[2k+1]\Gamma[5/2-k]} \right. \\ & \left. \times \left[ {}_2F_1\left(\frac{1}{2} - k, -k; \frac{5}{2} - k; \frac{1}{\omega^2}\right) - \left(k + \frac{3}{2}\right) {}_2F_1\left(\frac{3}{2} - k, -k; \frac{5}{2} - k; \frac{1}{\omega^2}\right) \right] \right] \end{aligned}$$

$$\begin{aligned} \mathcal{M}_{2\text{PM}}^{\text{odd}} = & -m_1 \left[ 4\mathcal{F}_1 + \sum_{k=0}^{\infty} \frac{(2\omega^2 - 1)}{(\omega^2 - 1)^{k+1}} \frac{(-1)^k 8\Gamma[k+1]}{\Gamma[2k+1]} \mathcal{F}_k V^k \right] \\ & - m_2 \left[ \frac{15\sqrt{\pi}}{2}\mathcal{F}_{1/2} + \sum_{k=0}^{\infty} \frac{\omega^{2k}}{(\omega^2 - 1)^{k+1}} \frac{4^{1-k} \mathcal{F}_k V^k}{(1)_k (2k-1)} \right. \\ & \left. \times \left[ {}_2F_1\left(-\frac{1}{2} - k, -k; \frac{3}{2} - k; \frac{1}{\omega^2}\right) - \left(k + \frac{5}{2}\right) {}_2F_1\left(\frac{1}{2} - k, -k; \frac{3}{2} - k; \frac{1}{\omega^2}\right) \right] \right] \end{aligned}$$

CoM dynamics if spin and angular momentum are aligned:



eikonal phase: 
$$\chi = \frac{1}{4m_1m_2\sqrt{\omega^2 - 1}} \int \frac{d^2\mathbf{q}}{(2\pi)^2} e^{i\mathbf{b}\cdot\mathbf{q}} \mathcal{M}_{2\text{PM}}$$

related simply to scattering angle [Amati, Ciafaloni, Veneziano '88]: 
$$\theta = -\frac{1}{|\mathbf{p}|} \frac{\partial}{\partial |\mathbf{b}|} \chi$$

$$\chi_{\text{aam}} = -\frac{G^2 \pi m_1 m_2}{4bx(1-x^2)^{3/2}(\omega^2-1)} \left( \frac{\chi_{\text{aam}}^{\text{even}}}{x\sqrt{(1-x^2)(\omega^2-1)}} + \omega \chi_{\text{aam}}^{\text{odd}} \right)$$

$$\chi_{\text{aam}}^{\text{even}} = -2m_1 \sqrt{1-x^2} \left[ x^2 \omega^2 (x^2 \omega^2 + 6(\omega^2 - 1)) + (1 - (1-x^2)^{3/2}) (\omega^2 - 1)^2 \right] \\ + m_2 x^2 \left[ 3(\omega^2 - 1) (5(x^2 - 1) \omega^2 - 2x^2 + 1) - 2x^2 \sqrt{1-x^2} \right]$$

$$\chi_{\text{aam}}^{\text{odd}} = 8m_1 \left[ \omega^2 x^2 + (\omega^2 - 1) (1 - (1-x^2)^{3/2}) \right] + 3x^2 m_2 \left[ 2 + 5(\omega^2 - 1) \sqrt{1-x^2} \right]$$

agrees with [Siemonsen, Vines '19] when  $\frac{m_2}{m_1} \rightarrow 0$ .



# Summary

Compton amplitude needed for 2PM Kerr  $\times$  Kerr scattering amplitude

known for Kerr up to fourth order in spin; unknown for higher spin, amplitude has spurious poles

derived viable classical amplitude to all spin orders; differs from Kerr only by contact terms

fixed contact terms assuming Kerr-scattering shift symmetry at low spins applies to all spins

spinning  $\times$  Schwarzschild amplitude derived to all spins

ultrarelativistic limit selects Kerr at fourth spin order; use limit to fix remaining coefficients for even-in-spin contact terms

**Kerr is not determined by this analysis... what is Kerr???**