

Unlocking the Inelastic Dark Matter window with Vector Mediators

Ana Luisa Foguel

in collaboration with Peter Reimitz and Renata Z. Funchal

based on: [arxiv:2410.00881](https://arxiv.org/abs/2410.00881)

J. High Energ. Phys. **2025**, 1 (2025)


afoguel@usp.br

University of São Paulo, São Paulo, Brazil

IFT/UAM, Madrid, Spain


DarkTools@Torino

16 June 2025





Outline

1. Introduction and Motivations
2. Inelastic Dark Matter
 - ↪ Theoretical framework
 - ↪ Decay Rates
3. Relic Density Computation
4. ReD-DeLiVeR code
5. Bounds
6. Conclusions




Introduction and Motivations





Introduction and Motivations

The Standard Model




DM ??





Introduction and Motivations

The Standard Model




DM ??

where is New Physics?
where is DM???



Introduction and Motivations



The Standard Model



DM ??


and what about...
neutrino masses?
BAU? Hierarchy problem?
Strong CP?

where is New Physics?
where is DM???




Introduction and Motivations

Where can we search for BSM signals?




Introduction and Motivations

Where can we search for BSM signals?




Introduction and Motivations

Where can we search for BSM signals?




Introduction and Motivations

Where can we search for BSM signals?




Introduction and Motivations

Where can we search for BSM signals?




Introduction and Motivations

How can we search for light particle BSM signals?




Introduction and Motivations

How can we search for light particle BSM signals?




Suppose we have particles χ and $\bar{\chi}$ in the dark sector

➤ natural possibility: couple to the gauge bosons of weak interactions




Introduction and Motivations

How can we search for light particle BSM signals?



Suppose we have particles χ and $\bar{\chi}$ in the dark sector

➔ natural possibility: couple to the gauge bosons of **weak interactions**




However...

$$\langle \sigma v \rangle \sim \frac{m_\chi^2}{m_Z^4}$$

which means that lowering the DM mass decreases the thermal-average cross section


➔ **DM is overproduced!**

Remember!




Introduction and Motivations

How can we search for light particle BSM signals?



Suppose we have particles χ and $\bar{\chi}$ in the dark sector

➔ natural possibility: couple to the gauge bosons of **weak interactions**



However...

$$\langle \sigma v \rangle \sim \frac{m_\chi^2}{m_Z^4}$$

which means that lowering the DM mass decreases the thermal-average cross section


→ **DM is overproduced!**

Lee-Weinberg bound

$$m_\chi \gtrsim 2 \text{ GeV}$$


Introduction and Motivations

How can we search for light particle BSM signals?



Introduction and Motivations

How can we search for light particle BSM signals?




➤ solution: inclusion of new light dark sector mediator states!

These light mediators will act as portals between the dark sector and the SM.



Introduction and Motivations • Hidden Portals

How can we search for light particle BSM signals?



Introduction and Motivations • Hidden Portals

How can we search for light particle BSM signals?



Inelastic Dark Matter

Theoretical Framework

Dark Sector

two-component Weyl fermions forming a Dirac pair

new vector mediator coming from a spontaneously broken $U(1)_Q$ abelian gauge symmetry


pseudo-Dirac pair

χ_1

χ_2



Inelastic Dark Matter (iDM)



From the diagonalization of the fermion mass terms

$$\mathcal{L} \supset -\overset{\text{Dirac}}{m_D \psi_1 \psi_2} - \overset{\text{Majorana}}{\frac{1}{2}(\delta_1 \psi_1^2 + \delta_2 \psi_2^2)} + \text{h.c.}, \quad \text{with} \quad \delta_{1,2} \ll m_D$$

$$\chi_1 \simeq \frac{i}{\sqrt{2}}(\psi_1 - \psi_2),$$

pseudo-Dirac pair with nearly degenerate masses

$$\chi_2 \simeq \frac{1}{\sqrt{2}}(\psi_1 + \psi_2),$$

$$m_{1,2} \simeq m_D \mp \frac{1}{2}(\delta_1 + \delta_2) \Rightarrow$$


Mass splitting

$$\Delta := \frac{m_2 - m_1}{m_1} = \frac{\delta_1 + \delta_2}{m_1} < 1$$

$\Rightarrow$ mass eigenstates

Inelastic Dark Matter

Theoretical Framework



From the diagonalization of the fermion mass terms

$$\mathcal{L} \supset -\overset{\text{Dirac}}{m_D \psi_1 \psi_2} - \overset{\text{Majorana}}{\frac{1}{2}(\delta_1 \psi_1^2 + \delta_2 \psi_2^2)} + \text{h.c.}, \quad \text{with} \quad \delta_{1,2} \ll m_D$$

$\Rightarrow$ mass eigenstates

$$\chi_1 \simeq \frac{i}{\sqrt{2}}(\psi_1 - \psi_2), \quad \text{pseudo-Dirac pair with nearly degenerate masses}$$

$$\chi_2 \simeq \frac{1}{\sqrt{2}}(\psi_1 + \psi_2), \quad m_{1,2} \simeq m_D \mp \frac{1}{2}(\delta_1 + \delta_2) \Rightarrow$$

Mass splitting


$$\Delta := \frac{m_2 - m_1}{m_1} = \frac{\delta_1 + \delta_2}{m_1} < 1$$

Inelastic Dark Matter

Theoretical Framework

The interaction term with the mediator turns to be **off-diagonal**

$$\mathcal{L} \supset g_D Z_{Q\mu} (\psi_1^\dagger \bar{\sigma}^\mu \psi_1 - \psi_2^\dagger \bar{\sigma}^\mu \psi_2) \longrightarrow \mathcal{L}_{\text{int}}^{\text{D}} = \frac{i}{2} g_D Z_{Q\mu} \bar{\chi}_2 \gamma^\mu \chi_1 + \text{h.c.}$$




Inelastic Dark Matter

Theoretical Framework

The interaction term with the mediator turns to be **off-diagonal**

$$\mathcal{L} \supset g_D Z_{Q\mu} (\psi_1^\dagger \bar{\sigma}^\mu \psi_1 - \psi_2^\dagger \bar{\sigma}^\mu \psi_2) \longrightarrow \mathcal{L}_{\text{int}}^{\text{D}} = \frac{i}{2} g_D Z_{Q\mu} \bar{\chi}_2 \gamma^\mu \chi_1 + \text{h.c.}$$



Motivations


➤ **Thermal relics:** DM abundance can be computed via thermal freeze-out.

Inelastic Dark Matter

Theoretical Framework

The interaction term with the mediator turns to be **off-diagonal**

$$\mathcal{L} \supset g_D Z_{Q\mu} (\psi_1^\dagger \bar{\sigma}^\mu \psi_1 - \psi_2^\dagger \bar{\sigma}^\mu \psi_2) \longrightarrow \mathcal{L}_{\text{int}}^{\text{D}} = \frac{i}{2} g_D Z_{Q\mu} \bar{\chi}_2 \gamma^\mu \chi_1 + \text{h.c.}$$



Motivations

➤ **Thermal relics:** DM abundance can be computed via thermal freeze-out.

➤ **Evades indirect and direct detection experimental limits**

The heavier state χ_2 can decay into the DM candidate χ_1 , depleting its abundance

⇒ **no present-day population of heavier states** to co-annihilate with the DM → avoid indirect detection signals


⇒ similarly, **direct detection** signals depend on **up-scatter** of the light state, which is **kinematically suppressed**

Inelastic Dark Matter

Theoretical Framework

The interaction term with the mediator turns to be **off-diagonal**

$$\mathcal{L} \supset g_D Z_{Q\mu} (\psi_1^\dagger \bar{\sigma}^\mu \psi_1 - \psi_2^\dagger \bar{\sigma}^\mu \psi_2) \longrightarrow \mathcal{L}_{\text{int}}^D = \frac{i}{2} g_D Z_{Q\mu} \bar{\chi}_2 \gamma^\mu \chi_1 + \text{h.c.}$$



Motivations

→ **Thermal relics:** DM abundance can be computed via thermal freeze-out.

→ **Evades indirect and direct detection experimental limits**

The heavier state χ_2 can decay into the DM candidate χ_1 , depleting its abundance

⇒ **no present-day population of heavier states** to co-annihilate with the DM → avoid indirect detection signals

⇒ similarly, **direct detection** signals depend on **up-scatter** of the light state, which is **kinematically suppressed**

→ **Evades stringent CMB limits**


Since the abundance of χ_2 is already reduced during recombination era, **coannihilations that would inject energy into the plasma are suppressed.**

Inelastic Dark Matter

Theoretical Framework



The interaction term with the mediator turns to be **off-diagonal**

$$\mathcal{L} \supset g_D Z_{Q\mu} (\psi_1^\dagger \bar{\sigma}^\mu \psi_1 - \psi_2^\dagger \bar{\sigma}^\mu \psi_2) \longrightarrow \mathcal{L}_{\text{int}}^D = \frac{i}{2} g_D Z_{Q\mu} \bar{\chi}_2 \gamma^\mu \chi_1 + \text{h.c.}$$



What's new?

- In the literature: only considered the minimal scenario with a secluded **dark photon portal** Z_D
However... this case has been nearly **completely ruled out by experimental limits**...



Inelastic Dark Matter

Theoretical Framework

The interaction term with the mediator turns to be **off-diagonal**

$$\mathcal{L} \supset g_D Z_{Q\mu} (\psi_1^\dagger \bar{\sigma}^\mu \psi_1 - \psi_2^\dagger \bar{\sigma}^\mu \psi_2) \longrightarrow \mathcal{L}_{\text{int}}^{\text{D}} = \frac{i}{2} g_D Z_{Q\mu} \bar{\chi}_2 \gamma^\mu \chi_1 + \text{h.c.}$$

What's new?


- This work: we consider the case of **generic charges for the U(1)_Q group**



vector mediator also couples to the SM via **direct terms** depending on the choice of charge

$$\mathcal{L}_{\text{int}}^{\text{SM}} = e \epsilon J_{\text{em}}^\mu Z_{Q\mu} - g_Q J_Q^\mu Z_{Q\mu}$$

$$J_Q^\mu = \sum_f q_Q^f \bar{f} \gamma^\mu f + \sum_{\ell=e,\mu,\tau} q_Q^{\nu_\ell} \bar{\nu}_\ell \gamma^\mu P_L \nu_\ell,$$



$$Q = x_B B - x_e L_e - x_\mu L_\mu - x_\tau L_\tau$$

x_B	x_e	x_μ	x_τ	Q	q_Q^f			
					quarks	e/ν_e	μ/ν_μ	τ/ν_τ
1	1	1	1	$B - L$	$\frac{1}{3}$	-1	-1	-1
1	0	0	3	$B - 3L_\tau$	$\frac{1}{3}$	0	0	-3
1	0	0	0	B	$\frac{1}{3}$	0	0	0
0	0	-1	1	$L_\mu - L_\tau$	0	0	1	-1

Inelastic Dark Matter

Theoretical Framework

The interaction term with the mediator turns to be **off-diagonal**

$$\mathcal{L} \supset g_D Z_{Q\mu} (\psi_1^\dagger \bar{\sigma}^\mu \psi_1 - \psi_2^\dagger \bar{\sigma}^\mu \psi_2) \longrightarrow \mathcal{L}_{\text{int}}^{\text{D}} = \frac{i}{2} g_D Z_{Q\mu} \bar{\chi}_2 \gamma^\mu \chi_1 + \text{h.c.}$$

What's new?

- This work: we consider the case of **generic charges for the $U(1)_Q$ group**

⇓

ALF, P. Reimitz, R.Z. Funchal
[arXiv:2410.00881](https://arxiv.org/abs/2410.00881) [hep-ph]


vector mediator also couples to the SM via **direct terms** depending on the choice of charge

$$\begin{aligned} \mathcal{L}_{\text{int}}^{\text{SM}} &= e \epsilon J_{\text{em}}^\mu Z_{Q\mu} - g_Q J_Q^\mu Z_{Q\mu} \\ J_Q^\mu &= \sum_f q_Q^f \bar{f} \gamma^\mu f + \sum_{\ell=e,\mu,\tau} q_Q^{\nu_\ell} \bar{\nu}_\ell \gamma^\mu P_L \nu_\ell, \end{aligned}$$

iDM_Q
models



free parameters: $m_{Z_Q}, R, \Delta, g_Q, \alpha_D$





$$Q = x_B B - x_e L_e - x_\mu L_\mu - x_\tau L_\tau$$

x_B	x_e	x_μ	x_τ	Q	q_Q^f			
					quarks	e/ν_e	μ/ν_μ	τ/ν_τ
1	1	1	1	$B - L$	$\frac{1}{3}$	-1	-1	-1
1	0	0	3	$B - 3L_\tau$	$\frac{1}{3}$	0	0	-3
1	0	0	0	B	$\frac{1}{3}$	0	0	0
0	0	-1	1	$L_\mu - L_\tau$	0	0	1	-1

Inelastic Dark Matter · Decay Rates

↪ Hierarchy $m_{Z_Q} > m_1 + m_2$
 ↪ Limit $g_D \gg g_Q$






Inelastic Dark Matter · Decay Rates

→ Hierarchy $m_{Z_Q} > m_1 + m_2$ → Limit $g_D \gg g_Q$

Dark fermion

$$\Gamma(\chi_2 \rightarrow \chi_1 \bar{f} f) \simeq \frac{4 \alpha_Q \alpha_D \Delta^5 m_{Z_Q}}{15 \pi R^5}$$



Inelastic Dark Matter · Relic Density Computation


Boltzmann Equation

$$\frac{dY_{1,2}}{dx} = \frac{s}{Hx} \left[-\langle\sigma v\rangle_{12\rightarrow ff} (Y_1 Y_2 - Y_1^{\text{eq}} Y_2^{\text{eq}}) \pm 2 \langle\sigma v\rangle_{22\rightarrow 11} \left((Y_2)^2 - \left(Y_1 \frac{Y_2^{\text{eq}}}{Y_1^{\text{eq}}} \right)^2 \right) \right. \\ \left. \pm \left(\langle\sigma v\rangle_{2f\rightarrow 1f} Y_f^{\text{eq}} + \frac{1}{s} \langle\Gamma\rangle_{2\rightarrow 1ff} \right) \left(Y_2 - Y_1 \frac{Y_2^{\text{eq}}}{Y_1^{\text{eq}}} \right) \right],$$

Inelastic Dark Matter · Relic Density Computation

Boltzmann Equation

$$\frac{dY_{1,2}}{dx} = \frac{s}{Hx} \left[-\langle\sigma v\rangle_{12\rightarrow ff} (Y_1 Y_2 - Y_1^{\text{eq}} Y_2^{\text{eq}}) \pm 2 \langle\sigma v\rangle_{22\rightarrow 11} \left((Y_2)^2 - \left(Y_1 \frac{Y_2^{\text{eq}}}{Y_1^{\text{eq}}} \right)^2 \right) \right. \\ \left. \pm \left(\langle\sigma v\rangle_{2f\rightarrow 1f} Y_f^{\text{eq}} + \frac{1}{s} \langle\Gamma\rangle_{2\rightarrow 1ff} \right) \left(Y_2 - Y_1 \frac{Y_2^{\text{eq}}}{Y_1^{\text{eq}}} \right) \right],$$




a) $\chi_1 \chi_2 \rightarrow \text{SM}$


Inelastic Dark Matter · Relic Density Computation

Boltzmann Equation

$$\frac{dY_{1,2}}{dx} = \frac{s}{Hx} \left[-\langle\sigma v\rangle_{12\rightarrow ff} (Y_1 Y_2 - Y_1^{\text{eq}} Y_2^{\text{eq}}) \pm 2\langle\sigma v\rangle_{22\rightarrow 11} \left((Y_2)^2 - \left(Y_1 \frac{Y_2^{\text{eq}}}{Y_1^{\text{eq}}} \right)^2 \right) \right. \\ \left. \pm \left(\langle\sigma v\rangle_{2f\rightarrow 1f} Y_f^{\text{eq}} + \frac{1}{s} \langle\Gamma\rangle_{2\rightarrow 1ff} \right) \left(Y_2 - Y_1 \frac{Y_2^{\text{eq}}}{Y_1^{\text{eq}}} \right) \right],$$



a) $\chi_1 \chi_2 \rightarrow \text{SM}$




b) $\chi_2 \chi_2 \rightarrow \chi_1 \chi_1$


Inelastic Dark Matter · Relic Density Computation

Boltzmann Equation


$$\frac{dY_{1,2}}{dx} = \frac{s}{Hx} \left[-\langle\sigma v\rangle_{12\rightarrow ff} (Y_1 Y_2 - Y_1^{\text{eq}} Y_2^{\text{eq}}) \pm 2\langle\sigma v\rangle_{22\rightarrow 11} \left((Y_2)^2 - \left(Y_1 \frac{Y_2^{\text{eq}}}{Y_1^{\text{eq}}} \right)^2 \right) \right. \\ \left. \pm \left(\langle\sigma v\rangle_{2f\rightarrow 1f} Y_f^{\text{eq}} + \frac{1}{s} \langle\Gamma\rangle_{2\rightarrow 1ff} \right) \left(Y_2 - Y_1 \frac{Y_2^{\text{eq}}}{Y_1^{\text{eq}}} \right) \right],$$



a) $\chi_1 \chi_2 \rightarrow \text{SM}$



b) $\chi_2 \chi_2 \rightarrow \chi_1 \chi_1$




c) $\chi_2 f \rightarrow \chi_1 f$


Inelastic Dark Matter · Relic Density Computation

Boltzmann Equation


$$\frac{dY_{1,2}}{dx} = \frac{s}{Hx} \left[-\langle\sigma v\rangle_{12\rightarrow ff} (Y_1 Y_2 - Y_1^{\text{eq}} Y_2^{\text{eq}}) \pm 2\langle\sigma v\rangle_{22\rightarrow 11} \left((Y_2)^2 - \left(Y_1 \frac{Y_2^{\text{eq}}}{Y_1^{\text{eq}}} \right)^2 \right) \right. \\ \left. \pm \left(\langle\sigma v\rangle_{2f\rightarrow 1f} Y_f^{\text{eq}} + \frac{1}{s} \langle\Gamma\rangle_{2\rightarrow 1ff} \right) \left(Y_2 - Y_1 \frac{Y_2^{\text{eq}}}{Y_1^{\text{eq}}} \right) \right],$$




a) $\chi_1 \chi_2 \rightarrow \text{SM}$



b) $\chi_2 \chi_2 \rightarrow \chi_1 \chi_1$



c) $\chi_2 f \rightarrow \chi_1 f$




d) $\chi_2 \rightarrow \chi_1 + \text{SM}$

Inelastic Dark Matter · Relic Density Computation


Boltzmann Equation

$$\frac{dY_{1,2}}{dx} = \frac{s}{Hx} \left[-\langle\sigma v\rangle_{12\rightarrow ff} (Y_1 Y_2 - Y_1^{\text{eq}} Y_2^{\text{eq}}) \pm 2\langle\sigma v\rangle_{22\rightarrow 11} \left((Y_2)^2 - \left(Y_1 \frac{Y_2^{\text{eq}}}{Y_1^{\text{eq}}} \right)^2 \right) \right. \\ \left. \pm \left(\langle\sigma v\rangle_{2f\rightarrow 1f} Y_f^{\text{eq}} + \frac{1}{s} \langle\Gamma\rangle_{2\rightarrow 1ff} \right) \left(Y_2 - Y_1 \frac{Y_2^{\text{eq}}}{Y_1^{\text{eq}}} \right) \right],$$


coannihilations
dominate!




a) $\chi_1 \chi_2 \rightarrow \text{SM}$



b) $\chi_2 \chi_2 \rightarrow \chi_1 \chi_1$



c) $\chi_2 f \rightarrow \chi_1 f$




d) $\chi_2 \rightarrow \chi_1 + \text{SM}$

Inelastic Dark Matter · Relic Density Computation

Boltzmann Equation

$$\frac{dY_{1,2}}{dx} = \frac{s}{Hx} \left[-\langle\sigma v\rangle_{12\rightarrow ff} (Y_1 Y_2 - Y_1^{\text{eq}} Y_2^{\text{eq}}) \pm 2 \langle\sigma v\rangle_{22\rightarrow 11} \left((Y_2)^2 - \left(Y_1 \frac{Y_2^{\text{eq}}}{Y_1^{\text{eq}}} \right)^2 \right) \right. \\ \left. \pm \left(\langle\sigma v\rangle_{2f\rightarrow 1f} Y_f^{\text{eq}} + \frac{1}{s} \langle\Gamma\rangle_{2\rightarrow 1ff} \right) \left(Y_2 - Y_1 \frac{Y_2^{\text{eq}}}{Y_1^{\text{eq}}} \right) \right],$$



we can simplify by considering


$$n = n_1 + n_2$$

$$\frac{dY}{dx} = -2 \frac{s}{xH} \langle\sigma v\rangle_{\text{eff}} (Y^2 - Y_{\text{eq}}^2)$$


$$\langle\sigma v\rangle_{\text{eff}} = \langle\sigma v\rangle_{12\rightarrow ff} \frac{n_1^{\text{eq}} n_2^{\text{eq}}}{(n^{\text{eq}})^2}$$

Inelastic Dark Matter · Relic Density Computation

iDM




iDM _{$L_\mu - L_\tau$}




Inelastic Dark Matter · Relic Density Computation

iDM



iDM_B



Inelastic Dark Matter · ReD-DeLiVeR code


ReD-DeLiVeR (Relic Density with DeLiVeR)

- update of the previous python package **DELIVER** (**Decays of Light Vectors Revised**), which can be used to compute decay rates and branching ratios for **user-defined $U(1)_Q$ charges**,
- **key feature**: includes a complete set of **hadronic decays** (20 channels)

channel	resonances
$\pi\gamma$	$\rho, \omega, \omega', \omega'', \phi$
$\pi\pi$	$\rho, \rho', \dots$
3π	$\rho, \rho'', \omega, \omega', \omega'', \phi$
4π	$\rho, \rho', \rho'', \rho'''$
KK	$\rho, \dots, \omega, \dots, \phi, \dots$
$KK\pi$	$\rho, \rho', \rho'', \phi, \phi', \phi''$

channel	resonances
$\eta\gamma$	$\rho, \rho', \omega, \phi$
$\eta\pi\pi$	ρ, ρ', ρ''
$\omega\pi \rightarrow \pi\pi\gamma$	ρ, ρ', ρ''
$\omega\pi\pi$	ω''
$\phi\pi$	ρ, ρ'
$\eta'\pi\pi$	ρ'''
$\eta\omega$	ω', ω''
$\eta\phi$	ϕ', ϕ''
$p\bar{p}/n\bar{n}$	$\rho, \rho', \dots, \omega, \omega', \dots$
$\phi\pi\pi$	ϕ', ϕ''
$K^*(892)K\pi$	ρ'', ϕ'
6π	ρ'''

ALF, P. Reimitz, R.Z. Funchal [JHEP 04 (2022)119]




Inelastic Dark Matter · ReD-DeLiVeR code

ReD-DeLiVeR (Relic Density with DeLiVeR)

- update of the previous python package **DELIVER** (**Decays of Light Vectors Revised**), which can be used to compute decay rates and branching ratios for **user-defined $U(1)_Q$ charges**,
- **key feature**: includes a complete set of **hadronic decays** (20 channels)
- **new version**:
 - inclusion of DM candidates
 - computation of **relic density** and **rates**

simplified DM models
inelastic DM




Inelastic Dark Matter · ReD-DeLiVeR code

ReD-DeLiVeR (Relic Density with DeLiVeR)

- update of the previous python package **DELIVER** (**Decays of Light Vectors Revised**), which can be used to compute decay rates and branching ratios for **user-defined $U(1)_Q$ charges**,
- **key feature**: includes a complete set of **hadronic decays** (20 channels)
- **new version**:
 - inclusion of DM candidates
 - computation of **relic density** and **rates**

simplified DM models
inelastic DM



Inelastic Dark Matter · ReD-DeLiVeR code


ReD-DeLiVeR (Relic Density with DeLiVeR)

→ update of the previous python package **DELIVER** (**D**ecays of **L**ight **V**ectors **R**evised), which can be used to compute decay rates and branching ratios for **user-defined $U(1)_Q$ charges**,

→ **key feature**: includes a complete set of **hadronic decays** (20 channels)

→ **new version**:

- inclusion of DM candidates
 - computation of **relic density** and **rates**
 - computation of **thermal targets**
- } simplified DM models
inelastic DM



specially for B-coupled models, it is very important to compute correctly the hadronic contributions

Inelastic Dark Matter · ReD-DeLiVeR code

ReD-DeLiVeR (Relic Density with DeLiVeR)

→ update of the previous python package **DELIVER** (**D**ecays of **L**ight **V**ectors **R**evised), which can be used to compute decay rates and branching ratios for **user-defined $U(1)_Q$ charges**,

→ **key feature**: includes a complete set of **hadronic decays** (20 channels)

→ **new version**:

- inclusion of DM candidates
 - computation of **relic density** and **rates**
 - computation of **thermal targets**
- } simplified DM models
inelastic DM


→ **publicly available** on GitHub together with a **Tutorial!**

<https://github.com/anafoguel/ReD-DeLiVeR>

ReD-DeLiVeR

by Ana Luisa Foguel, Peter Reimitz, and Renata Zukanovich Funchal



arXiv: 2410.00881



specially for B-coupled models, it is very important to compute correctly the hadronic contributions

Inelastic Dark Matter · Bounds



→ $i\text{DM}_{B-L}$



Inelastic Dark Matter · Bounds




iDM_B




Inelastic Dark Matter · Bounds

→ **iDM_B**





⇒





Inelastic Dark Matter · Bounds

→ $i\text{DM}_{B-3L_\tau}$





Inelastic Dark Matter · Bounds

→ $i\text{DM}_{B-3L_\tau}$



Inelastic Dark Matter · Bounds

→ iDM $_{L_\mu - L_\tau}$



Conclusions

- **Light Feebly Interacting Particles** can shed light in several unanswered questions of the SM
- As experiments increase their **luminosities**, and we enter the **intensity frontier** era of particle physics, we increase the capabilities to probe new light sectors.
- As a guiding principle, we consider different **portals** between the **Dark Sector** and the **SM**
- In this work we considered a **vector portal** to a fermionic **inelastic Dark Matter** sector

thermal relics evade indirect and direct detection evade CMB bounds

general vector mediators

$i\text{DM}_Q$

- We developed a code that computes the relic density **ReD-DeLiVeR**
- With general mediators, we showed that we can **unlock new regions of the parameter space** of the vanilla dark photon model

$$\curvearrowright B - 3L_\tau \quad \curvearrowright L_\mu - L_\tau$$

Conclusions

- **Light Feebly Interacting Particles** can shed light in several unanswered questions of the SM
- As experiments increase their **luminosities**, and we enter the **intensity frontier** era of particle physics, we increase the capabilities to probe new light sectors.
- As a guiding principle, we consider different **portals** between the **Dark Sector** and the **SM**
- In this work we considered a **vector portal** to a fermionic **inelastic Dark Matter** sector

thermal relics evade indirect and direct detection evade CMB bounds

general vector mediators

iDM_Q

- We developed a code that computes the relic density **ReD-DeLiVeR**
- With general mediators, we showed that we can **unlock new regions of the parameter space** of the vanilla dark photon model

$$\curvearrowright B - 3L_\tau \quad \curvearrowright L_\mu - L_\tau$$

*Thank you for your
kind attention!*

Thank you
for your
attention!!



BACKUP

Inelastic Dark Matter

DM Thermal Freeze-out · WIMP miracle

We know that freeze-out happens when $\Gamma \sim H$

$$m_\chi \sim \alpha_{\text{eff}} \sqrt{T_{\text{eq}} M_{\text{Pl}}} \sim \alpha_{\text{eff}} \times 30 \text{ TeV}$$


WIMP miracle

↪ For couplings similar to the electroweak coupling ($\alpha_{\text{eff}} \sim 10^{-2}$) $\Rightarrow$ EW scale emerges naturally


However, this also implies that

$$m_{\text{DM}} \gtrsim \frac{m_Z^2}{(T_{\text{eq}} m_{\text{Pl}})^{1/2}} \sim \text{GeV}.$$

Hence, sub-GeV DM motivates the presence of new light mediators



Inelastic Dark Matter






arXiv:2311.00032v1 [hep-ph]

Inelastic Dark Matter · Decay Rates

Decay rates · Mediator

→ Hierarchy $m_{Z_Q} > m_1 + m_2$

→ Limit $g_D \gg g_Q$






prompt-decay

Inelastic Dark Matter · Decay Rates


Decay rates · Dark fermion χ_2





$$\Gamma(\chi_2 \rightarrow \chi_1 \bar{f} f) \simeq \frac{4 \alpha_Q \alpha_D \Delta^5 m_{Z_Q}}{15 \pi R^5}$$





Inelastic Dark Matter · Decay Rates



Decay rates · Dark fermion χ_2









Simulation Checks





Different thermal targets



Fixed Dark Charges

$$\mathcal{L}_{\text{int}}^{\text{D}} = i g_D Z_{Q\mu} \bar{\chi}_2 \gamma^\mu \chi_1 + \text{h.c.}$$

$$g_D \equiv g_Q q_D$$



Details on the Experimental Limits

- Invisible searches

BaBar, Belle II, LDMX, NA62 and NA64-e

$$\frac{\sigma_{Z_Q}}{\sigma_{Z_\gamma}} \{ \text{BR}(Z_Q \rightarrow \chi_1 \chi_2) [\text{BR}(\chi_2 \rightarrow \chi_1 \bar{\nu} \nu) + P_{\text{out}}^{\chi_2} \text{BR}(\chi_2 \rightarrow \chi_1 + \text{vis})] + \text{BR}(Z_Q \rightarrow \bar{\nu} \nu) \} - 1 = 0$$

$$P_{\text{out}}^{\chi_2} = e^{-L_{\text{out}}/\lambda_{\chi_2}}$$

↪ recast from invisible Dark photon limits

- Semi-visible searches

E137, CHARM, NuCal and LSND

$$\frac{\sigma_{Z_Q}}{\sigma_{Z_\gamma}} \left(\frac{\text{BR}(Z_Q \rightarrow \chi_1 \chi_2)}{\text{BR}(Z_\gamma \rightarrow \chi_1 \chi_2)} \right) \left(\frac{\text{BR}(\chi_2 \rightarrow Z_Q^* \rightarrow \chi_1 e^+ e^-)}{\text{BR}(\chi_2 \rightarrow Z_\gamma^* \rightarrow \chi_1 e^+ e^-)} \right) \left(\frac{\varepsilon_Q(\tau_{\chi_2})}{\varepsilon_\gamma(\tau_{\chi_2})} \right) - 1 = 0$$

↪ recast from the original iDM limits

Details on the Experimental Limits

- 4μ searches from BaBar and CMS

↪ constrained $L_\mu - L_\tau$ mediators,

$$\frac{g_Q}{g_{L_\mu - L_\tau}^{\text{exp}}} \left[\frac{\text{BR}(Z_Q \rightarrow \mu^+ \mu^-) + \text{BR}(Z_Q \rightarrow \chi_1 \chi_2) \text{BR}(\chi_2 \rightarrow \chi_1 \mu^+ \mu^-) P_{\text{in}}^{\chi_2}}{\text{BR}(Z_{L_\mu - L_\tau} \rightarrow \mu^+ \mu^-)} \right] - 1 = 0$$

$$P_{\text{in}}^{\chi_2} = e^{-L_{\text{in}}/\lambda_{\chi_2}} - e^{-L_{\text{out}}/\lambda_{\chi_2}}$$

- Meson decays

J/Ψ and Υ invisible decay searches

$$\begin{aligned} \frac{\text{BR}(J/\Psi \rightarrow \chi_1 \chi_2)}{\text{BR}(J/\Psi \rightarrow e^+ e^-)} &= \frac{\alpha_D}{\alpha_e} \left(\frac{q_Q^c g_Q}{q_{\text{em}}^c e} \right)^2 \frac{m_V^4}{(m_{Z_Q}^2 - m_V^2)^2 + m_{Z_Q}^2 \Gamma_{Z_Q}^2} \\ &\times \left(1 - \frac{\Delta^2}{R^2} m_r^2 \right)^{3/2} \left(1 + \frac{m_r^2 (\Delta + 2)^2}{2R^2} \right) \sqrt{1 - \frac{m_r^2 (\Delta + 2)^2}{R^2}} \end{aligned}$$

$$\begin{aligned} \text{BR}(V \rightarrow \text{inv})|_{\text{iDM}_Q} &= \text{BR}(V \rightarrow \chi_1 \chi_2) [\text{BR}(\chi_2 \rightarrow \chi_1 \bar{\nu} \nu) + P_{\text{out}}^{\chi_2} \text{BR}(\chi_2 \rightarrow \chi_1 + \text{vis})] \\ &\quad + \text{BR}(V \rightarrow Z_Q^* \rightarrow \bar{\nu} \nu). \end{aligned} \quad \Rightarrow \quad \text{BR}(V \rightarrow \text{inv})|_{\text{iDM}_Q} < \text{BR}(V \rightarrow \text{inv})|_{\text{exp}}$$